

REVIEW

Circulating vitamin D level and mortality in prostate cancer patients: a dose–response meta-analysis

Zhen-yu Song¹, Qiuming Yao², Zhiyuan Zhuo¹, Zhe Ma¹ and Gang Chen¹

¹Department of Urology, Jinshan Hospital of Fudan University, Shanghai, China

²Department of Endocrinology, Jinshan Hospital of Fudan University, Shanghai, China

Correspondence should be addressed to G Chen: chgan305@163.com

Abstract

Previous studies investigating the association of circulating 25-hydroxyvitamin D level with prognosis of prostate cancer yielded controversial results. We conducted a dose–response meta-analysis to elucidate the relationship. PubMed and EMBASE were searched for eligible studies up to July 15, 2018. We performed a dose–response meta-analysis using random-effect model to calculate the summary hazard ratio (HR) and 95% CI of mortality in patients with prostate cancer. Seven eligible cohort studies with 7808 participants were included. The results indicated that higher vitamin D level could reduce the risk of death among prostate cancer patients. The summary HR of prostate cancer-specific mortality correlated with an increment of every 20 nmol/L in circulating vitamin D level was 0.91, with 95% CI 0.87–0.97, $P=0.002$. The HR for all-cause mortality with the increase of 20 nmol/L vitamin D was 0.91 (95% CI: 0.84–0.98, $P=0.01$). Sensitivity analysis suggested the pooled HRs were stable and not obviously changed by any single study. No evidence of publications bias was observed. This meta-analysis suggested that higher 25-hydroxyvitamin D level was associated with a reduction of mortality in prostate cancer patients and vitamin D is an important protective factor in the progression and prognosis of prostate cancer.

Key Words

- ▶ vitamin D
- ▶ mortality
- ▶ prostate cancer
- ▶ meta-analysis

Endocrine Connections
(2018) 7, R294–R303

Introduction

Prostate cancer (PCa) is one of the most common malignant tumors in male. In 2017, American Cancer Society reported 161,360 cases of newly diagnosed PCa, accounting for 20% of male tumors. Furthermore, its incidence and mortality ranked the first place and third respectively (1). The mortality of PCa was proposed to be associated with obesity, physical activity, smoking, antioxidants, etc. (2). At present, the treatment of PCa have caused serious economic burden (3). More useful treatment measures are urgently needed by people to improve the survival rate of prostate cancer patients.

The major circulating form of vitamin D in human body is 25-hydroxyvitamin D (25(OH)D), which comes

from vitamin D via 25-hydroxylation process in the liver. 25(OH)D can be converted into 1,25(OH)₂D by 1 α -hydroxylase, which is the most active hormonal metabolite of vitamin D. As a hormone, 1,25(OH)₂D binds to vitamin D receptor located in nucleus and functions. It is reported to play an important role in cellular proliferation (4), differentiation, apoptosis (5), angiogenesis (6) and metastasis (7). All these processes may regulate the development and progression of cancer.

A number of researches have been done to clarify the association between vitamin D and PCa. Some experimental studies indicated that vitamin D might play a crucial role in the occurrence and progression of

PCa. One study demonstrated mutations of vitamin D receptor gene were associated with Gleason score (8). Furthermore, study showed that genetic variants in the vitamin D pathway had effects on the risk of progression, prostate cancer-specific mortality and recurrence of PCa (9). Recent studies have reported controversial results about the association of vitamin D with the survival rate of prostate cancer. For example, in newly diagnosed stage IV prostate cancer patients, no significant association of 25-hydroxyvitamin D with the prognosis of them was found (10). In contrast, other studies reported that higher 25-hydroxyvitamin D was related to improved prostate cancer prognosis (11, 12).

Therefore, the relationship between 25-hydroxyvitamin D level and mortality of PCa is still unclear. Hence, we conducted this analysis to explore whether circulating 25-hydroxyvitamin D level was correlated with the survival of PCa through a dose-response meta-analysis.

Materials and methods

Search strategy

We searched PubMed and EMBASE databases from inception to July 15, 2018, for eligible studies on the relationship between vitamin D and mortality in prostate cancer patients. The terms used to retrieve literatures were the following: (vitamin D OR 25-hydroxyvitamin D OR 25(OH)D) and (prostate cancer OR prostate carcinoma). We also referred to the reference lists from reviews or relevant papers to get more eligible researches. There was no language restriction.

Selection criteria

Reports were included in this dose-response meta-analysis if they met the criteria as follows: (1) the association between vitamin D and mortality in prostate cancer patients was reported; (2) the study type was cohort; (3) the risk estimates of mortality in prostate cancer patients, like HR and 95% CI were reported. If the same data were used in several studies, we selected the publication with the largest number of cases or more details.

Data extraction

Data were extracted from eligible studies by two researchers independently. The information collected

from each study contained of the first author's last name, publication year, country, follow-up time, number of cases and person-year, risk estimates with corresponding 95% CIs and confounding factors adjusted in multivariable analysis. We extracted the risk estimates from the most completed adjusted model to decrease the risk of possible confounding. Disagreements were resolved by consensus among authors.

Quality assessment

We evaluated the quality of studies by use of the Newcastle Ottawa Scale (NOS) (13). According to its criteria, studies were assessed on the basis of three perspectives: selection, comparability and outcomes. If studies got seven or more stars, they were regarded as high quality. Differences were resolved by discussion.

Statistical analysis

We performed data analyses separately for two outcomes, namely all-cause mortality and prostate cancer-specific mortality. Pooled HRs were calculated to assess the impact of vitamin D level on the prognosis of patients. The method proposed by Greenland and Longnecker (14) and Orsini *et al.* (15) was used to estimate the HR per 20 nmol/L




Figure 1 Flowchart of study selection in the meta-analysis.

Table 1 The main characteristics of the included studies in the meta-analysis.

| Study | Country | Study design | Time of vitamin D assessment | Participants | Follow-up | Outcomes | Age at diagnosis (years) | Adjustments | Quality |
|------------------------------------|---------|--------------------|------------------------------|--------------|------------|-----------|--------------------------|---|---------|
| Tretli <i>et al.</i> 2009 (23) | Norway | Cohort | Postdiagnosis | 160 | 44 months | ACM; PCSM | 64.5 | Patient group and age, tumor differentiation grade and the patient functional status at the time of blood collection | 7 |
| Fang <i>et al.</i> 2011 (11) | USA | Prospective cohort | Prediagnosis | 1822 | 10 years | ACM; PCSM | 68.9 | Age at diagnosis, body mass index, physical activity, and smoking, Gleason score, and TNM stage | 9 |
| Holt <i>et al.</i> 2013 (35) | USA | Prospective cohort | Postdiagnosis | 1476 | 10.8 years | PCSM | 60 | Season of blood draw, age and race, BMI, smoking status, and weekly exercise stage, Gleason score and primary treatment | 9 |
| Gupta <i>et al.</i> 2015 (10) | USA | Prospective cohort | Postdiagnosis | 125 | 31 months | ACM; PCSM | 60 | Age, ECOG performance status, BMI, prostate specific antigen (PSA), season of blood draw, CTCA hospital, serum albumin, corrected serum calcium, bone metastasis and nutritional status | 7 |
| Mondul <i>et al.</i> 2016 (28) | Finland | Prospective cohort | Prediagnosis | 1000 | 23 years | PCSM | 69.2 | Age, physical activity, cigarettes per day, and family history of prostate cancer | 9 |
| Meyer <i>et al.</i> 2016 (32) | Norway | Prospective cohort | Prediagnosis | 2282 | 21.2 years | ACM | NA | Age, month of blood sampling and examination physical activity, BMI, smoking and education | 9 |
| Brandstedt <i>et al.</i> 2016 (34) | Sweden | Prospective cohort | Prediagnosis | 943 | 16.6 years | ACM; PCSM | 69.3 | Season and year of inclusion, age at baseline, age at diagnosis, body mass index (BMI), and tumor characteristics (TNM and Gleason score) | 9 |

ACM, all-cause mortality; BMI, body mass index; CTCA, Cancer Treatment Centers of America; ECOG, Eastern Cooperative Oncology Group; NA, not available; PCSM, prostate cancer-specific mortality; PSA, prostate specific antigen.

increase of vitamin D level. Statistical heterogeneity among studies was evaluated with the use of Q and I^2 statistic (16, 17). For the Q statistic, we regarded P value <0.10 as statistically significant heterogeneity among studies. As to the I^2 statistic, I^2 more than 50% also suggested obvious heterogeneity. We utilized the random-effects model to combine HRs from single studies if obvious heterogeneity was observed (18). In the sensitivity analysis, studies were omitted one by one and the others were analyzed to evaluate the effect of single study on the summary risk estimates. Publication bias was assessed with the use of funnel plot and the Egger's test (19). We utilized Stata (Version 12.0) to perform this dose–response analysis. P value <0.05 was reckoned as statistically significant difference.

Results

Study selection and characteristics

The selection process was showed in Fig. 1. We retrieved 2650 articles from PubMed and EMBASE databases (Fig. 1). A majority of them were excluded from our analysis because they did not belong to cohort studies or because outcomes were not associated with our analysis, leaving 19 articles for detailed evaluation by reading full-texts (10, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37). Twelve studies were then removed after reading their full-texts. Two studies were excluded because of inadequate study design (22, 24). Nine studies were excluded because they did not contain prognosis data among prostate cancer patients (20, 21, 26, 27, 29, 32, 35, 37). One study was not qualified as a result of unusable data (36). Finally, a total of seven studies were included into our meta-analysis. The seven studies were published between 2009 and 2016 and the total number of prostate cancer participants was 7808. All of them were performed in developed countries, written in English (Table 1). Among them, three studies were conducted in USA (10, 30, 34), two in Norway (23, 31), one in Finland (28), one in Sweden (33). All studies were prospective cohort type, except one from Tretli S. It is also a cohort study but hard to define it belongs to prospective or retrospective type. Meanwhile, the vitamin D assessments were performed after diagnosis in three studies, while the others were before diagnosis of prostate cancer. All studies reported adjusted HRs. Every research was adjusted for many confounding factors, such as age, BMI, drinking history and so forth. Participants were followed up from 4 to 21 years. Five studies contained HRs of all-cause mortality among prostate cancer patients, and six reported HRs of prostate cancer-specific mortality.

The quality assessment of those studies according to NOS criteria was also presented in the Table 1.

25-hydroxyvitamin D and all-cause mortality

We observed significant heterogeneity among five studies on all-cause mortality ($I^2=68.9\%$). Figure 2A displayed the results of the dose–response analyses on all-cause mortality (Fig. 2A). A nonlinear relationship existed between 25-hydroxyvitamin D and risk of all-cause mortality in prostate cancer patients, suggesting higher 25-hydroxyvitamin D level was associated with decreased risk of death from all causes among prostate cancer patients ($P=0.038$). The summary HR of all-cause mortality correlated with an increment of every 20 nmol/L in circulating vitamin D level was 0.91 (95% CI:




Figure 2

Dose–response relationships between 25(OH)D and risk estimates of all-cause mortality and prostate cancer-specific mortality. (A) Risk estimates with 95% CI for the association between 25(OH)D and all-cause mortality. (B) Risk estimates with 95% CI for the association between 25(OH)D and prostate cancer-specific mortality.

0.84–0.98, $P=0.01$) (Fig. 3A). Sensitivity analysis suggested the pooled HRs were stable and not obviously changed by any individual study (Fig. 4A).

25-hydroxyvitamin D and prostate cancer-specific mortality

There was obvious heterogeneity observed among those six studies on prostate cancer-specific mortality ($I^2=53.4%$). A nonlinear relationship between 25-hydroxyvitamin D and risk of prostate cancer-specific mortality was also presented in Fig. 2B, indicating higher vitamin D level could decrease the mortality from prostate cancer (Fig. 2B). The summary HR of prostate cancer-specific mortality correlated with an increment of every 20nmol/L in circulating vitamin D level were 0.91 (95% CI: 0.87–0.97, $P=0.002$) (Fig. 3B). The sensitivity analysis showed the summary HRs were

not markedly changed by any individual study (Fig. 4B), indicating no significant influence of single study on the results.

Publication bias

No risk of publication bias was observed in the funnel plots (Fig. 5). The outcomes from Egger's test also suggested that there were no publication bias for the analysis of all-cause mortality ($P=0.143$) and prostate cancer-specific mortality ($P=0.301$).

Subgroup analysis and meta-regression

We conducted the subgroup analysis and meta-regression to detect the source of heterogeneity, which was presented in Table 2. Stratifying by the time of vitamin D assessment,




Figure 3 Summary risk estimates of mortality in prostate cancer patients associated with 20nmol/L increment in 25(OH)D level. (A) Funnel plot of risk estimates of all-cause mortality of prostate cancer with the increment of 20nmol/L in 25(OH)D level. (B) Funnel plot of risk estimates of prostate cancer-specific mortality with the increment of 20nmol/L in 25(OH)D level.




Figure 4 Sensitivity analysis by excluding studies by turns suggested that the pooled HRs were not significantly changed by any individual study. (A) Sensitivity analysis of the association between 25(OH)D and all-cause mortality of prostate cancer. (B) Sensitivity analysis of the association between 25(OH)D and prostate cancer-specific mortality.

the HR of prostate cancer-specific mortality was 0.91 (95% CI: 0.88–0.95) for prediagnosis studies and 0.84 (95% CI: 0.58–1.21) for postdiagnosis ones. The HR of all-cause mortality was 0.94 (95% CI: 0.88–0.98) in prediagnosis subgroup. Restricting the analysis among more than 10-year follow-up yielded a HR of 0.92 (95% CI: 0.89–0.96) and 0.94 (95% CI: 0.89–0.98) for prostate cancer-specific mortality and all-cause mortality respectively, which was slightly higher than the overall results. Moreover, there was no evidence of significant heterogeneity between subgroups with the use of meta-regression analyses.

Discussion

The role of circulating 25-hydroxyvitamin D and survival outcomes among prostate cancer patients remains unclear

and controversial. This meta-analysis is the first one to focus on the relationship between 25-hydroxyvitamin D and mortality in prostate cancer, involving 7808 participants with survival outcomes. The results calculated from seven eligible studies indicated higher vitamin D level was significantly associated with decreased all-cause mortality and prostate cancer-specific mortality. Further dose-response analysis showed that every 20 nmol/L increment in 25-hydroxyvitamin D level was associated with a 9% lower risk of all-cause mortality and prostate cancer-specific mortality. By conducting the subgroup analysis, we found the results were consistent in prediagnosis and more than 10-year follow-up subgroups. The assessment of vitamin D before diagnosis was more likely to get rid of the influence of prostate cancer on the level of vitamin D and long follow-up time enabled researchers to calculate the outcome events more precisely. Based on the above findings, we conclude that higher circulating vitamin D




Figure 5 Publication bias. (A) Publication bias of the association between 25(OH)D and all-cause mortality of prostate cancer. (B) Publication bias of the association between 25(OH)D and prostate cancer-specific mortality.

Table 2 Summary risk estimates of the associations between vitamin D level and prostate cancer mortality.

| Study characteristics | No. of studies | HR | 95% CI | I ² (%) | P value 1 | P value 2 |
|------------------------------|----------------|------|-----------|--------------------|-----------|-----------|
| Studies of PCM | 6 | 0.91 | 0.87–0.97 | 53.4 | 0.057 | |
| Country | | | | | | 0.294 |
| Europe | 4 | 0.88 | 0.81–0.95 | 57.9 | 0.068 | |
| USA | 2 | 0.96 | 0.90–1.03 | 0 | 0.389 | |
| Time of vitamin D assessment | | | | | | 0.36 |
| Postdiagnosis | 2 | 0.84 | 0.58–1.21 | 89.1 | 0.002 | |
| Prediagnosis | 4 | 0.91 | 0.88–0.95 | 0 | 0.675 | |
| Follow-up | | | | | | 0.055 |
| Less than 10 years | 1 | | | | | |
| More than 10 years | 5 | 0.92 | 0.89–0.96 | 0 | 0.479 | |
| Studies of ACM | 5 | 0.91 | 0.84–0.98 | 68.9 | 0.012 | |
| Country | | | | | | 0.295 |
| Europe | 3 | 0.87 | 0.79 | 68.5 | 0.042 | |
| USA | 2 | 0.98 | 0.93–1.03 | 0 | 0.576 | |
| Time of vitamin D assessment | | | | | | 0.246 |
| Postdiagnosis | 2 | 0.83 | 0.66–1.04 | 71.5 | 0.061 | |
| Prediagnosis | 3 | 0.94 | 0.89–0.98 | 53.9 | 0.114 | |
| Follow-up | | | | | | 0.246 |
| Less than 10 years | 2 | 0.83 | 0.66–1.04 | 71.5 | 0.061 | |
| More than 10 years | 3 | 0.94 | 0.89–0.98 | 53.9 | 0.114 | |

P value 1 for heterogeneity within each subgroup. P value 2 for heterogeneity between subgroups with meta-regression analysis. ACM, all-cause mortality; CI, confidence interval; HR, summary hazard ratio; PCSM, prostate cancer-specific mortality.

level is associated with a lower risk of death from prostate cancer.

Numerous experimental studies have been done to elucidate the mechanism by which vitamin D affect the prostate cancer survival. According to previous studies, 1,25(OH)₂D could cause cell cycle arrest and induce apoptosis, inhibiting cell proliferation in several prostate cancer cell lines (38, 39, 40). 1,25(OH)₂D played a protective role in preventing normal human prostate epithelial cell lines from oxidative stress in since it increased both the expression and activity of antioxidants, such as glucose-6-phosphate dehydrogenase and glutathione (41). Ben-Shoshan and colleagues demonstrated that 1,25(OH)₂D inhibited angiogenesis by reducing HIF-1 α expression in various human prostate cancer cell lines (42). In terms of animal model evidence, Ray and colleagues indicated that a diet deficient in vitamin D rather than vitamin D-sufficient diet accelerated growth of human prostate cancers insensitive to androgen therapy in athymic mice (43). Another study reported that a higher vitamin D3-supplemented diet led to significant tumor shrinkage in mice bearing PC-3 prostate cancer xenografts (44). Moreover, vitamin D could prevent the metastasis of prostate cancer according to several animal and cell experiments (45, 46). Therefore, there is some evidence supporting the protective effect of vitamin D in prostate cancer. However, the underlying molecular mechanisms

are still not fully clarified, and more studies are needed to explore them.

Some studies reported that 25-hydroxyvitamin D concentration was correlated with prostate cancer pathology. Researchers found lower 25-hydroxyvitamin D concentrations were positively correlated with higher Gleason grade and tumor stage (47, 48). The findings above provide some explanations for the prognostic role of 25-hydroxyvitamin D in prostate cancer.

Previous studies reported conflicting results about the vitamin D and prostate cancer incidence. One meta-analysis showed positive association between high level of vitamin D and increased incidence of prostate cancer (49). Some studies also suggested that high incidence of aggressive prostate cancer in African Americans might be partly due to deficient concentrations of serum vitamin D (50, 51). In the contrast, one Mendelian randomization study showed null relationship between vitamin D and risk of prostate cancer (52). Other studies also failed to find a positive relationship between vitamin D and prostate cancer risk (47, 53). The conflicting findings in the relationship between vitamin D and prostate cancer risk may result from the some factors, such as different populations, various study design and different confounding factors. The findings in our study suggest that vitamin D is more likely to be a suppressive and protective factor during the development of prostate

cancer. Therefore, there is still controversy on the role of vitamin D in prostate cancer, which need to be elucidated in future researches.

There is also some evidence from clinical trials on the roles of vitamin D in prostate cancer. In a clinical trial, low-grade prostate cancer patients took 4000IU of vitamin D3 every day for a whole year and had a biopsy after the supplementation (54). Results of biopsy revealed a decreased number of positive cores and no increase in Gleason score (54). Several randomized clinical trials showed that oral vitamin D3 modestly decreased the level of PSA (55) and reduced the PSA rise rate (56, 57). However, a vitamin D supplementation trial showed no influence on free or total PSA level in African American population (58). At present, the evidence from clinical trials on the roles of vitamin D in prostate cancer is still limited, and more clinical trials are needed.

There are potential limitations existing in our study which should be considered. For one thing, although all studies adjusted for confounding factors, some potential confounding factors related to vitamin D remained residual. For another, some studies included in our meta-analysis tested the circulating vitamin D level postdiagnosis or post treatment, thus it is difficult to get rid of the possibility of reverse causality. What is more, the limited number of included studies restricted us to find the source of heterogeneity.

Based on the results mentioned earlier, we can draw the conclusion that higher vitamin D level is significantly associated with a risk reduction of all-cause mortality and prostate cancer-specific mortality, indicating vitamin D may exert a protective effect in the progression and prognosis of prostate cancer. More cohort studies and randomized clinical trial are needed to further illustrate the role of vitamin D in the pathogenesis and prognosis of prostate cancer.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

Author contribution statement

Zhen-yu Song designed the study. Qiuming Yao, Zhi-yuan Zhuo and Zhe Ma extracted the data. Zhen-yu Song and Qiuming Yao performed the analyses. Zhen-yu Song wrote the draft. Gang Chen revised it critically.

References

- 1 Siegel RL, Miller KD & Jemal A. Cancer statistics, 2017. *CA: A Cancer Journal for Clinicians* 2017 **67** 7–30. (<https://doi.org/10.3322/caac.21387>)
- 2 Wilson KM, Giovannucci EL & Mucci LA. Lifestyle and dietary factors in the prevention of lethal prostate cancer. *Asian Journal of Andrology* 2012 **14** 365–374. (<https://doi.org/10.1038/aja.2011.142>)
- 3 Crawford ED, Black L, Eaddy M & Kruep EJ. A retrospective analysis illustrating the substantial clinical and economic burden of prostate cancer. *Prostate Cancer and Prostatic Diseases* 2010 **13** 162–167. (<https://doi.org/10.1038/pcan.2009.63>)
- 4 Scaglione-Sewell BA, Bissonnette M, Skarosi S, Abraham C & Brasitus TA. A vitamin D3 analog induces a G1-phase arrest in CaCo-2 cells by inhibiting cdk2 and cdk6: roles of cyclin E, p21Waf1, and p27Kip1. *Endocrinology* 2000 **141** 3931–3939. (<https://doi.org/10.1210/endo.141.11.7782>)
- 5 Diaz GD, Paraskeva C, Thomas MG, Binderup L & Hague A. Apoptosis is induced by the active metabolite of vitamin D3 and its analogue EB1089 in colorectal adenoma and carcinoma cells: possible implications for prevention and therapy. *Cancer Research* 2000 **60** 2304–2312.
- 6 Fernandez-Garcia NI, Palmer HG, Garcia M, Gonzalez-Martin A, Del RM, Barettino D, Volpert O, Munoz A & Jimenez B. 1alpha,25-Dihydroxyvitamin D3 regulates the expression of Id1 and Id2 genes and the angiogenic phenotype of human colon carcinoma cells. *Oncogene* 2005 **24** 6533–6544. (<https://doi.org/10.1038/sj.onc.1208801>)
- 7 Evans SR, Shchepotin EI, Young H, Rochon J, Uskokovic M & Shchepotin IB. 1,25-dihydroxyvitamin D3 synthetic analogs inhibit spontaneous metastases in a 1,2-dimethylhydrazine-induced colon carcinogenesis model. *International Journal of Oncology* 2000 **16** 1249–1254. (<https://doi.org/10.3892/ijo.16.6.1249>)
- 8 Chen L, Davey SG, Evans DM, Cox A, Lawlor DA, Donovan J, Yuan W, Day IN, Martin RM, Lane A, *et al.* Genetic variants in the vitamin d receptor are associated with advanced prostate cancer at diagnosis: findings from the prostate testing for cancer and treatment study and a systematic review. *Cancer Epidemiology, Biomarkers and Prevention* 2009 **18** 2874–2881. (<https://doi.org/10.1158/1055-9965.EPI-09-0544>)
- 9 Holt SK, Kwon EM, Koopmeiners JS, Lin DW, Feng Z, Ostrander EA, Peters U & Stanford JL. Vitamin D pathway gene variants and prostate cancer prognosis. *Prostate* 2010 **70** 1448–1460. (<https://doi.org/10.1002/pros.21180>)
- 10 Gupta D, Trukova K, Popiel B, Lammersfeld C & Vashi PG. The association between pre-treatment serum 25-hydroxyvitamin D and survival in newly diagnosed stage IV prostate cancer. *PLoS ONE* 2015 **10** e119690. (<https://doi.org/10.1371/journal.pone.0119690>)
- 11 Fang F, Kasperzyk JL, Shui I, Hendrickson W, Hollis BW, Fall K, Ma J, Gaziano JM, Stampfer MJ, Mucci LA & Giovannucci E. Prediagnostic plasma vitamin D metabolites and mortality among patients with prostate cancer. *PLoS ONE* 2011 **6** e18625. (<https://doi.org/10.1371/journal.pone.0018625>)
- 12 Brandstedt J, Almquist M, Manjer J & Malm J. Vitamin D, PTH, and calcium in relation to survival following prostate cancer. *Cancer Causes and Control* 2016 **27** 669–677. (<https://doi.org/10.1007/s10552-016-0740-7>)
- 13 Margulis AV, Pladevall M, Riera-Guardia N, Varas-Lorenzo C, Hazell L, Berkman ND, Viswanathan M & Perez-Gutthann S. Quality assessment of observational studies in a drug-safety systematic review, comparison of two tools: the Newcastle-Ottawa Scale and the RTI item bank. *Clinical Epidemiology* 2014 **6** 359–368. (<https://doi.org/10.2147/CLEP.S66677>)
- 14 Greenland S & Longnecker MP. Methods for trend estimation from summarized dose-response data, with applications to meta-analysis.

- American Journal of Epidemiology* 1992 **135** 1301–1309. (<https://doi.org/10.1093/oxfordjournals.aje.a116237>)
- 15 Orsini N, Bellocco R & Greenland S. Generalized least squares for trend estimation of summarized dose–response data. *Stata Journal* 2006 **6** 40–57.
 - 16 Higgins JP, Thompson SG, Deeks JJ & Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003 **327** 557–560. (<https://doi.org/10.1136/bmj.327.7414.557>)
 - 17 Cochran WG. The combination of estimates from different experiments. *Biometrics* 1954 **10** 101–129. (<https://doi.org/10.2307/3001666>)
 - 18 DerSimonian R & Laird N. Meta-analysis in clinical trials. *Controlled Clinical Trials* 1986 **7** 177–188. ([https://doi.org/10.1016/0197-2456\(86\)90046-2](https://doi.org/10.1016/0197-2456(86)90046-2))
 - 19 Egger M, Davey SG, Schneider M & Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997 **315** 629–634. (<https://doi.org/10.1136/bmj.315.7109.629>)
 - 20 Feldman D, Krishnan AV, Swami S, Giovannucci E & Feldman BJ. The role of vitamin D in reducing cancer risk and progression. *Nature Reviews Cancer* 2014 **14** 342–357. (<https://doi.org/10.1038/nrc3691>)
 - 21 Giovannucci E, Liu Y, Rimm EB, Hollis BW, Fuchs CS, Stampfer MJ & Willett WC. Prospective study of predictors of vitamin D status and cancer incidence and mortality in men. *Journal of the National Cancer Institute* 2006 **98** 451–459. (<https://doi.org/10.1093/jnci/djj101>)
 - 22 Shui IM, Mucci LA, Kraft P, Tamimi RM, Lindstrom S, Penney KL, Nimptsch K, Hollis BW, Dupre N, Platz EA, *et al.* Vitamin D-related genetic variation, plasma vitamin D, and risk of lethal prostate cancer: a prospective nested case-control study. *Journal of the National Cancer Institute* 2012 **104** 690–699. (<https://doi.org/10.1093/jnci/djs189>)
 - 23 Tretli S, Hernes E, Berg JP, Hestvik UE & Robsahm TE. Association between serum 25(OH)D and death from prostate cancer. *British Journal of Cancer* 2009 **100** 450–454. (<https://doi.org/10.1038/sj.bjc.6604865>)
 - 24 Shui IM, Mondul AM, Lindstrom S, Tsilidis KK, Travis RC, Gerke T, Albanes D, Mucci LA, Giovannucci E & Kraft P. Circulating vitamin D, vitamin D-related genetic variation, and risk of fatal prostate cancer in the National Cancer Institute Breast and Prostate Cancer Cohort Consortium. *Cancer* 2015 **121** 1949–1956. (<https://doi.org/10.1002/cncr.29320>)
 - 25 Buttigliero C, Monagheddu C, Petroni P, Saini A, Dogliotti L, Ciccone G & Berruti A. Prognostic role of vitamin d status and efficacy of vitamin D supplementation in cancer patients: a systematic review. *Oncologist* 2011 **16** 1215–1227. (<https://doi.org/10.1634/theoncologist.2011-0098>)
 - 26 Dimitrakopoulou VI, Travis RC, Shui IM, Mondul A, Albanes D, Virtamo J, Agudo A, Boeing H, Bueno-de-Mesquita HB, Gunter MJ, *et al.* Interactions between genome-wide significant genetic variants and circulating concentrations of 25-hydroxyvitamin D in relation to prostate cancer risk in the National Cancer Institute BPC3. *American Journal of Epidemiology* 2017 **185** 452–464. (<https://doi.org/10.1093/aje/kww143>)
 - 27 Porojnicu A, Robsahm TE, Berg JP & Moan J. Season of diagnosis is a predictor of cancer survival. Sun-induced vitamin D may be involved: a possible role of sun-induced Vitamin D. *Journal of Steroid Biochemistry and Molecular Biology* 2007 **103** 675–678. (<https://doi.org/10.1016/j.jsbmb.2006.12.031>)
 - 28 Mondul AM, Weinstein SJ, Moy KA, Mannisto S & Albanes D. Circulating 25-hydroxyvitamin D and prostate cancer survival. *Cancer Epidemiology, Biomarkers and Prevention* 2016 **25** 665–669. (<https://doi.org/10.1158/1055-9965.EPI-15-0991>)
 - 29 Paller CJ, Kanaan YM, Beyene DA, Naab TJ, Copeland RL, Tsai HL, Kanarek NF & Hudson TS. Risk of prostate cancer in African-American men: evidence of mixed effects of dietary quercetin by serum vitamin D status. *Prostate* 2015 **75** 1376–1383. (<https://doi.org/10.1002/pros.23018>)
 - 30 Fang F, Kasperzyk JL, Shui I, Hendrickson W, Hollis BW, Fall K, Ma J, Gaziano JM, Stampfer MJ, Mucci LA, *et al.* Prediagnostic plasma vitamin D metabolites and mortality among patients with prostate cancer. *PLoS ONE* 2011 **6** e18625. (<https://doi.org/10.1371/journal.pone.0018625>)
 - 31 Meyer HE, Stoer NC, Samuelsen SO, Blomhoff R, Robsahm TE, Brustad M, Giovannucci EL & Bjorge T. Long term association between serum 25-hydroxyvitamin D and mortality in a cohort of 4379 men. *PLoS ONE* 2016 **11** e151441. (<https://doi.org/10.1371/journal.pone.0151441>)
 - 32 Robsahm TE, Tretli S, Dahlback A & Moan J. Vitamin D3 from sunlight may improve the prognosis of breast-, colon- and prostate cancer (Norway). *Cancer Causes and Control* 2004 **15** 149–158. (<https://doi.org/10.1023/B:CACO.0000019494.34403.09>)
 - 33 Brandstedt J, Almquist M, Manjer J & Malm J. Vitamin D, PTH, and calcium in relation to survival following prostate cancer. *Cancer Causes and Control* 2016 **27** 669–677. (<https://doi.org/10.1007/s10552-016-0740-7>)
 - 34 Holt SK, Kolb S, Fu R, Horst R, Feng Z & Stanford JL. Circulating levels of 25-hydroxyvitamin D and prostate cancer prognosis. *Cancer Epidemiology* 2013 **37** 666–670. (<https://doi.org/10.1016/j.canep.2013.07.005>)
 - 35 Grant WB. The likely role of vitamin D from solar ultraviolet-B irradiance in increasing cancer survival. *Anticancer Research* 2006 **26** 2605–2614.
 - 36 Der T, Bailey BA, Youssef D, Manning T, Grant WB & Peiris AN. Vitamin D and prostate cancer survival in veterans. *Military Medicine* 2014 **179** 81–84. (<https://doi.org/10.7205/MILMED-D-12-00540>)
 - 37 Atoum MF, AlKateeb D & AlHaj MS. The Fok1 vitamin D receptor gene polymorphism and 25(OH) D serum levels and prostate cancer among Jordanian men. *Asian Pacific Journal of Cancer Prevention* 2015 **16** 2227–2230. (<https://doi.org/10.7314/APJCP.2015.16.6.2227>)
 - 38 Blut SE, McDonnell TJ, Polek TC & Weigel NL. Calcitriol-induced apoptosis in LNCaP cells is blocked by overexpression of Bcl-2. *Endocrinology* 2000 **141** 10–17. (<https://doi.org/10.1210/endo.141.1.7289>)
 - 39 Ikeda N, Uemura H, Ishiguro H, Hori M, Hosaka M, Kyo S, Miyamoto K, Takeda E & Kubota Y. Combination treatment with 1alpha,25-dihydroxyvitamin D3 and 9-cis-retinoic acid directly inhibits human telomerase reverse transcriptase transcription in prostate cancer cells. *Molecular Cancer Therapeutics* 2003 **2** 739–746.
 - 40 Rohan JN & Weigel NL. 1Alpha,25-dihydroxyvitamin D3 reduces c-Myc expression, inhibiting proliferation and causing G1 accumulation in C4-2 prostate cancer cells. *Endocrinology* 2009 **150** 2046–2054. (<https://doi.org/10.1210/en.2008-1395>)
 - 41 Bao BY, Ting HJ, Hsu JW & Le YF. Protective role of 1 alpha, 25-dihydroxyvitamin D3 against oxidative stress in nonmalignant human prostate epithelial cells. *International Journal of Cancer* 2008 **122** 2699–2706. (<https://doi.org/10.1002/ijc.23460>)
 - 42 Ben-Shoshan M, Amir S, Dang DT, Dang LH, Weisman Y & Mabjeesh NJ. 1alpha,25-dihydroxyvitamin D3 (Calcitriol) inhibits hypoxia-inducible factor-1/vascular endothelial growth factor pathway in human cancer cells. *Molecular Cancer Therapeutics* 2007 **6** 1433–1439. (<https://doi.org/10.1158/1535-7163.MCT-06-0677>)
 - 43 Ray R, Banks M, Abuzahra H, Eddy VJ, Persons KS, Lucia MS, Lambert JR & Holick MF. Effect of dietary vitamin D and calcium on the growth of androgen-insensitive human prostate tumor in a murine model. *Anticancer Research* 2012 **32** 727–731.
 - 44 Swami S, Krishnan AV, Wang JY, Jensen K, Horst R, Albertelli MA & Feldman D. Dietary vitamin D(3) and 1,25-dihydroxyvitamin D(3) (calcitriol) exhibit equivalent anticancer activity in mouse xenograft models of breast and prostate cancer. *Endocrinology* 2012 **153** 2576–2587. (<https://doi.org/10.1210/en.2011-1600>)
 - 45 Krishnan AV, Shinghal R, Raghavachari N, Brooks JD & Peehl DM. Analysis of vitamin D-regulated gene expression in LNCaP human

- prostate cancer cells using cDNA microarrays. *Prostate* 2004 **59** 243–251. (<https://doi.org/10.1002/pros.20006>)
- 46 Tokar EJ & Webber MM. Cholecalciferol (vitamin D3) inhibits growth and invasion by up-regulating nuclear receptors and 25-hydroxylase (CYP27A1) in human prostate cancer cells. *Clinical and Experimental Metastasis* 2005 **22** 275–284. (<https://doi.org/10.1007/s10585-005-8393-z>)
- 47 Gilbert R, Metcalfe C, Fraser WD, Donovan J, Hamdy F, Neal DE, Lane JA & Martin RM. Associations of circulating 25-hydroxyvitamin D with prostate cancer diagnosis, stage and grade. *International Journal of Cancer* 2012 **131** 1187–1196. (<https://doi.org/10.1002/ijc.27327>)
- 48 Nyame YA, Murphy AB, Bowen DK, Jordan G, Batai K, Dixon M, Hollowel CM, Kielb S, Meeks JJ, Gan PH, *et al.* Associations between serum vitamin D and adverse pathology in men undergoing radical prostatectomy. *Journal of Clinical Oncology* 2016 **34** 1345–1349. (<https://doi.org/10.1200/JCO.2015.65.1463>)
- 49 Xu Y, Shao X, Yao Y, Xu L, Chang L, Jiang Z & Lin Z. Positive association between circulating 25-hydroxyvitamin D levels and prostate cancer risk: new findings from an updated meta-analysis. *Journal of Cancer Research and Clinical Oncology* 2014 **140** 1465–1477. (<https://doi.org/10.1007/s00432-014-1706-3>)
- 50 Nelson SM, Batai K, Ahaghotu C, Agurs-Collins T & Kittles RA. Association between serum 25-hydroxy-vitamin D and aggressive prostate cancer in African American men. *Nutrients* 2016 **9** 12. (<https://doi.org/10.3390/nu9010012>)
- 51 Jackson MD, Tulloch-Reid MK, Lindsay CM, Smith G, Bennet FI, McFarlane-Anderson N, Aiken W & Coard KC. Both serum 25-hydroxyvitamin D and calcium levels may increase the risk of incident prostate cancer in Caribbean men of African ancestry. *Cancer Medicine* 2015 **4** 925–935. (<https://doi.org/10.1002/cam4.457>)
- 52 Dimitrakopoulou VI, Tsilidis KK, Haycock PC, Dimou NL, Al-Dabhani K, Martin RM, Lewis SJ, Gunter MJ, Mondul A, Shui IM, *et al.* Circulating vitamin D concentration and risk of seven cancers: mendelian randomisation study. *BMJ* 2017 **359** j4761. (<https://doi.org/10.1136/bmj.j4761>)
- 53 Ordóñez-Mena JM, Schöttker B, Fedirko V, Jenab M, Olsen A, Halkjær J, Kampman E, de Groot L, Jansen E, Bueno-de-Mesquita HB, *et al.* Pre-diagnostic vitamin D concentrations and cancer risks in older individuals: an analysis of cohorts participating in the CHANCES consortium. *European Journal of Epidemiology* 2016 **31** 311–323. (<https://doi.org/10.1007/s10654-015-0040-7>)
- 54 Marshal DT, Savage SJ, Garrett-Mayer E, Keane TE, Hollis BW, Horst RL, Ambrose LH, Kindy MS & Gattoni-Celli S. Vitamin D3 supplementation at 4000 international units per day for one year results in a decrease of positive cores at repeat biopsy in subjects with low-risk prostate cancer under active surveillance. *Journal of Clinical Endocrinology and Metabolism* 2012 **97** 2315–2324. (<https://doi.org/10.1210/jc.2012-1451>)
- 55 Wagner D, Trudel D, Van der Kwast T, Nonn L, Giangreco AA, Li D, Dias A, Cardoza M, Laszlo S, Hersey K, Klotz L, Finelli A, Fleshner N & Vieth R. Randomized clinical trial of vitamin D3 doses on prostatic vitamin D metabolite levels and ki67 labeling in prostate cancer patients. *Journal of Clinical Endocrinology and Metabolism* 2013 **98** 1498–1507. (<https://doi.org/10.1210/jc.2012-4019>)
- 56 Gross C, Stamey T, Hancock S & Feldman D. Treatment of early recurrent prostate cancer with 1,25-dihydroxyvitamin D3 (calcitriol). *Journal of Urology* 1998 **159** 2035–2039 2039–2040. ([https://doi.org/10.1016/S0022-5347\(01\)63236-1](https://doi.org/10.1016/S0022-5347(01)63236-1))
- 57 Beer TM, Lemmon D, Lowe BA & Henner WD. High-dose weekly oral calcitriol in patients with a rising PSA after prostatectomy or radiation for prostate carcinoma. *Cancer* 2003 **97** 1217–1224. (<https://doi.org/10.1002/cncr.11179>)
- 58 Chandler PD, Giovannucci EL, Scot JB, Bennet GG, Ng K, Chan AT, Hollis BW, Emmons KM, Fuchs CS & Drake BF. Null association between vitamin D and PSA levels among black men in a vitamin D supplementation trial. *Cancer Epidemiology, Biomarkers and Prevention* 2014 **23** 1944–1947. (<https://doi.org/10.1158/1055-9965.EPI-14-0522>)

 Received in final form 3 September 2018

Accepted 8 October 2018

Accepted Preprint published online 9 October 2018